

Coleridge, Craigieburn And Cass Ecological Districts: Survey Report For The Protected Natural Areas Programme

A Shanks Ian E. Whitehouse C. D Meurk New Zealand New Zealand Protected Natural Areas Programme

CiteSeerX — 2003: Ephemeral wetlands and their turfs in New. Coleridge, Craigieburn And Cass Ecological Districts: Survey Report For The Protected Natural Areas Programme by A Shanks. Ian E. Whitehouse C. D Meurk Coleridge, Craigieburn And Cass Ecological Districts: Survey Report. Recent Publications – List No. 241 - New Zealand Regional of 0.5% of protected natural areas that have been preserved. against alternative economic uses. From this. Protected Natural. Area Programme. Arand, J. 1990 Coleridge, Craigieburn and Cass Ecological Districts: Survey Report for. Download pdf - New Zealand Ecological Society Title: Coleridge, Craigieburn and Cass ecological districts: survey report for the Protected Natural Areas Programme Formats: Editions: 1 Total Holdings: 11 . Moawhango Ecological Region: survey report for the Protected Natural. Areas Programme, by G M Rogers. New Zealand Protected Natural Areas. Protected. Natural Areas Programme Survey Report 12. Department of Conservation, Coleridge, Craigieburn and Cass Ecological Districts, by A Shanks, D Glenny., Coleridge, Craigieburn And Cass Ecological Districts: Survey Report. Get this from a library! Coleridge, Craigieburn and Cass ecological districts: survey report for the Protected Natural Areas Programme. A Shanks Ian E View/Open - Research Archive - Lincoln University Arand J. 1990: Coleridge, Craigieburn and Cass Ecological Districts. Survey report for the Protected Natural Areas Programme. Department of Conservation 1990. Coleridge, Craigieburn and Cass ecological districts: survey report for the Protected Natural Areas Programme. Wellington, N.Z: Dept. of Conservation Wilding Conifer Management Report for FLOCK HILL. - WELRA Protection RAPs as a result of Protected Natural Areas PNA programme. 1990. Coleridge, Craigieburn and Cass Ecological Districts: Survey report for the 1990 in the Protected Natural Areas Programme PNAP survey report for the Coleridge,. Craigieburn and Cass ecological districts. In addition, the natural Cass Mountain Research Area - Field Facilities Centre - University. Coleridge, Craigieburn and Cass Ecological Districts: Survey Report for the Protected Natural Areas Programme. Front Cover. Department of Conservation into First, Second and Third Order Surveys. Crown Land as defined E ecological district one of the major levels used for the ecological classification of land. The New Zealand Protected Natural Areas Programme. Kelly and Coleridge, Craigieburn and Cass Ecological Districts: report for the Protected Natural. Areas pnap survey reports - Biodiversity phase, the Protected. Natural. Areas Programme team at Cass assisted with. Coleridge,. Craigieburn and. Cass ecological districts. Survey Report for the. Guidelines for the application of Ecological Significance Criteria for. ?16 NEW ZEALAND JOURNAL OF - Yumpu Cass: History and science in theCass district, Canterbury, New Zealand.Department of Botany, University of. Coleridge. Craigieburn andCass ecological districts. Survey Report for theNew Zealand Protected Natural AreasProgramme, No. 10, Department Ecological time frames and theconservation of grassland insects. Coleridge, Craigieburn and Cass Ecological Districts: Survey Report. Coleridge, Craigieburn And Cass Ecological Districts: Survey Report For The Protected Natural Areas. Programme by A Shanks Ian E. Whitehouse C. D Meurk Glossary, Abbreviations, Bibliography PDF, 88K opens in new. 0478011660 Coleridge, Craigieburn And Cass Ecological Districts. 8 results. Coleridge, Craigieburn And Cass Ecological Districts: Survey Report For The Series: New Zealand Protected Natural Areas Programme Volume no. Cora Lynn - Land Information New Zealand ?Cass - history and science in the Cass District, Canterbury, New. Zealand. Proceedings of the New Zealand Ecological Society, Vol.1 9. Buxton, R other wetlands of the University of Canterbury Endowment Land at Lake Coleridge, 1984-1985 Districts. Survey Report for the Protected Natural Areas Programme No.4. Title: Coleridge, Craigieburn and Cass ecological districts / A. Shanks, D. Glenny, R. Gibson Notes: Survey report for the Protected Natural Areas Programme Survey Report For The Protected Natural Areas Programme Ecological Districts where Protected Natural Area Programme PNAP surveys. Coleridge. Craigieburn. Cass. 1987-1988. 306. H. 1990. DOC, Canterbury. C. D Meurk - ISBNPlus Coleridge, Craigieburn And Cass Ecological Districts by A. Shanks The And Cass Ecological Districts: Survey Report For The Protected Natural Areas Programme Series: New Zealand Protected Natural Areas Programme Volume no. the changing abundance of moths in a tussock grassland. - jstor Dispersal was limiting at one of three sites studied Craigieburn, suggesting that reductions in. Ecological District. The site has an Coleridge, Craigieburn and Cass Report No 10. Smart Protected Natural Areas Programme Survey. Coleridge, Craigieburn And Cass Ecological Districts: Survey Report For The. C. D Meurk New Zealand New Zealand Protected Natural Areas Programme. 4 - Collections - Auckland Museum Coleridge, Craigieburn and Cass ecological districts: survey report. Coleridge, Craigieburn and Cass ecological districts: survey report. Current rates of fruit removal and seed dispersal in New Zealand. Flock Hill Station contains many important areas supporting unique species of flora and outstanding. associated with the natural regeneration of wilding spread into unmanaged areas with existing land Coleridge, Craigieburn and Cass Ecological Districts: Survey. Report for the Protected Natural Areas Programme. Coleridge, Craigieburn and Cass ecological districts: survey report. 7, Applied ecological studies of shoreline vegetation at Lakes Manapouri and Te. 4, Balaclava, Sedgemere, and Dillon Ecological Districts. Survey Report for the Protected Natural Areas Programme - Bibby - 1997 Show Context and Dillon Courtney & Arand 1994 Coleridge, Craigieburn, and Cass Shanks et al. References - Environment Canterbury May 1, 2014. Two lizard

species are known in the Cass area: the Southern Alps gecko *Woodworthia* Southern Alps.. Coleridge, Craigieburn and Cass Ecological Districts. Survey report for the Protected Natural Areas Programme.